

## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <a href="http://about.jstor.org/participate-jstor/individuals/early-journal-content">http://about.jstor.org/participate-jstor/individuals/early-journal-content</a>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

## INDEX TO VOLUME XIV\*

New names and the final members of new combinations are in **bold face type** 

Abies, 177, 181, 182; grandis, 174, 179, 180, 182, 183; pectinata, 8, 10 Absidia glauca, 149; septata, 149 Acer, 5, 6, 9, 181; circinnatum, 175, 198; macrophyllum, 174, 182, 183, 192; spicatum, 238 Achillea lanulosa, 118; Millefolium, Achlya, 152, 154, 155, 166; prolifera, Addition to the distribution of a rare fungus, 49 Aecidium, 105, 114; arcularium, 106; Pourthiaeae, 293, 295 Agarics, Dark-spored — I, 61; — II, 121; — III, 200; — IV, 258 Agaricus, 27, 75, 125, 136, 200, 220, 269; abruptibulbus, 209; abruptus, 209; Achimenes, 220; acuminatus, 128; adnatus, 133; aeruginosus, 128; alabamensis, 202; amygdalinus, 220, 221; appendiculatus, 64; approximans, 215; argenteus, 211; Artemisiae, 75; arvensis, 208, 210; atomatus, 275; atrorufus, 263; brunnescens, 211; bulbosus, 209; bulbularis, 131; bullaceus, 263; camarophyllus atramentosus, 48; camarophyllus fuligineus, 48; campester, 136, 203, 205, 212; campester hortensis, 212; campestris, 136; campestris sylvicola, 209; Candolleanus, 71; capnoides, 74; chlamydopus, 205; Comaropsis, 75; comtuliformis, 203; comtulus, 202, 203; coprophilus, 259; cornilla, 131; coronatus, 71; cothurnatus, 205; cretacellus, 208, 221; cretaceus, 220; depilatus, 134; diminutivus, 189, 214; disseminatus, 1270; distans, 126; echinatus, 213; elixus, 48; eludens, 217; eximius, 97; fabaceus, 220, 221;

falcifolius, 278; fascicularis, 73; Feildeni, 141; flavescens, 27, 221; floridanus, 207; foederatus, 221; glutinosus, 122, 133; graciloides, 277; haemorrhoidarius, 215, 216, 219; halophilus, 212; hiascens, 278; hirtosquamulosus, 75; Hornemanni, 135; Howeanus, 141; hymenocephalus, 64; hypoxanthus, 66; incertus, 64; Johnsonianus, 141; lacrymabundus, 67; lateritius, 72, 73; maculatus, 124; madeodiscus, 62; magniceps, 213; magnificus, 213; maritimus, 212; melanospermus, 131; melaspermus, 130, 131; merdarius, 138; micromegethus, 210; minutulus, 270; nitidipes, 75; obturatus, 141; odoratus, 276; olivaesporus, 97; ornellus, 75; oxyosmus, 214; perplexus, 72; pholidotus, 268; pilosporus, 204; placomyces, 217, 218; polytrichophilus, 261; populinus, 66; praerimosus, 204; pratensis, 208; pusillus, 210; Rodmani, 207; rubribrunnescens, 216; rupincola, 278; rutilescens, 211; saccharinophilus, 64; semiglobatus, 132; semivestitus, 269; solidipes, 203; sphaerosporus, 204; squamosus, 126; stercorarius, 133; Sterlingii, 217; Storea, 66; subcernuus, 126; sublateritius, 72; subrufescens, 189, 206, 210, 219, 221; sylvaticus, 218; sylvicola, 209, 221; tabularis, 204; thraustus aurantiacus, 127; tricholepis, 76; umbonatescens, 128; variabilis, 206; velutinus, 67; violaceolamellatus, 71; viridulus, 128; viscidus, 122; xylogenus, 221

Agropyron divergens, 279; occidentale, 279; repens, 279; Smithii, 279; spicatum, 279

\* It has been considered unnecessary to include here the species listed in the three following articles, since they are already indexed or specially listed.

Arthur: Uredinales collected by Fred J. Seaver in Trinidad. See p. 23, 24. Murrill: Index to illustrations of fungi, XXIII-XXXIII. See p. 332.

Wann and Muenscher: A preliminary list of the myxomycetes of the Cayuga Lake Basin. See p. 39.

Agrostis alba, 279; palustris, 279

Aira caespitosa, 33, 34

Aleuria aurantia, 175

Aleurodiscus, 179; amorphus, 179; subcruentatus, 179 Allescheria, 242; Boydii, 242 Alnus, 2, 5, 174, 175 Alopecurus aristulatus, 229, 230 Alternaria, 337 Amanita, 185; calyptrata, 185; chlorinosma, 185; muscaria, 185, 315; pantherina, 92; pantherinoides, 185; phalloides, 317; solitaria, 185; spissa, 92 Amanitopsis plumbea, 315; vaginata, 185; vaginata alba, 186; vaginata livida, 186 Amauroderma, 42; Gusmanianum, 43; Mosselmanii, 43; picipes, 43 Amelanchier, 2, 102, 289; asiatica, 288, 294; Cusickii, 91 American porias - I. Diagnoses of, 1 Amphisphaeria diffusa, 237; subiculosa, 237 Amsonia ciliata, 252, 297 Anellaria, 130; fimiputris, 130 Anemone, 176 Anisocarpua, 115 Another green-spored genus of gillfungi, 96 Aplospora, 336; Nyssae, 336 Aposphaeria brunneotincta, 102 Arachnion album, 195 Arcangeliella caudata, 196 Aristida, 230 Armillaria, 187; albolanaripes, 186; alphitophylla, 338; corticata, 187; nardosmia, 25 Arnica, 114; cana, 114; foliosa, 114, Artemisia, 118; tridentata, 45 Arthur, J. C., Uredinales collected by Fred J. Seaver in Trinidad, 12 Articles, Notes and brief, 42, 90, 231, Arundinaria Simoni, 86 Ascomycete isolated from a granular mycetoma of man, Life history of an undescribed, 239 Ascomycetes — I, Studies in tropical, Ascophora mucedo, 143 Aster, 302, 306-310; acuminatus, 303, 309; Chapmanii, 304; concinnus, 309; conspicuus, 254, 297, 303-306; cordifolius, 246, 254, 303-306; corrigiatus, 309; divaricatus, 303, 306; ericoides. dumosus, 303, 304; 254, 303; Fremontii, 309; fron-

dosus.

309;

Geyeri,

303-306;

hesperius, 254, 303; hirsuticaulis, 309; junceus, 309; laevigatus, 303; laevis, 246, 254, 303-306; laevis Geyeri, 254; lentus, 254, 303, 304; longifolius, 297, 306; Lowrieanus, 303, 309; macrophyllus, 303-306; novi-belgii, 304; oblongifolius, 309; paniculatus, 254, 303, 304, 306; patens, 303; patulus, 309; Pringlei, 306, 309; puniceus, 303; salicifolius, 303; schistosus, 309; spectabilis, 309; tenuicaulis, 309; undulatus, 246, 254, 303-306; vimineus, 303, 304, 306; viscosum, 309 Atylospora, 258, 262, 264, 273, 278; australis, 266, 278; corrugis, 267, 268; microsperma, 264; multipedata, 265; prunuliformis, 266, 278; striatula, 267, 278; umbonata, 267, 268; vestita, 265 Auricularia nigrescens, 44

Bacidia inundata, 95 Bacterium marginatum, 336 Balsamorrhiza, 112; macrophylla, 112 Barroetia, 107 Battarrea, 45; laciniata, 45 Betula, 5 Boehmeria nivea, 88 Boletinus cavipes, 181 Boletus, 45; chrysenteron, 180; luridus, 180; luteus, 25, 45, 180, 198; scaber, 180; scaber fuligineus, 180; scaber testaceus, 180; Zelleri, 181 Borrichia arborescens, 111; frutescens, III Botrydium, 95 Botryodiplodia Theobromae, 91 Bovista, 321; montana, 193; plumbea, 193 Bremia, 88 Brickellia, 107 Bromus, 279; ciliatus, 279 Broteria, 42 Bulgaria inquinans, 175 Bullaria, 105, 111, 112; Kuhniae, 106 Byssosphaeria diffusa, 237

Caeomusus triannulatus, 111
Calamagrostis canadensis, 279
Calendula, 120; officinalis, 119
Calicium Curtisii, 103; **Rhois**, 103
Callistephus chinensis, 306
Calocera cornea, 178
Calonyction, 299; aculeatum, 297—299; grandiflorum, 298
Calosphaeria, 100
Calvatia gigantea, 193; lilacina, 193; lilacina occidentalis, 193; rubroflava, 193

Calycadenia, 115 Camarophyllus, 48 Camillea Leprieurii, 338 Campanula rapunculoides, 305 Campanularius, 130 Cantharellus aurantiacus, 188; cibarius, 317; floccosus, 188; infundibuliformis, 188 Carduaceae, New or noteworthy rusts 011, 104 Carex, 104, 177; pendula, 33 Carpinus, 1 Carya illinoensis, 77 Castanea dentata, 102 Catastoma circumscissum, 193 Cayuga Lake Basin, Myxomycetes of the, 38 Cenangium balsameum, 101; querneum, 101; turgidum, 101 Centromadia, 115 Cephalosporium, 242; Boydii, 242 Cercospora Brunkii, 91; fusimaculans, 198; minima, 89; piricola, 88; tortipes, 337; tuberculella, 337 Cerotetium Dicentrae, 336 Chamaecyparis, 285; pisifera, 294; pisifera plumosa, 294; pisifera squarrosa, 294 Chamonixia, 196, 197; caespitosa, 197 Chanterel, 96; floccosus, 25, 43 Characters of the southern tuckahoe, Chloroneuron, 96 Chlorophyllum, 96 Chlorosperma, 96; olivaespora, 97 Chrysopsis Mariana, 246, 248, 298, 303-306 Cinnamomum camphora, 83 Circinella, 166; conica, 149, 160-162; minor, 161-163 Cladochytrium, 84; Maydis, 84 Cladoderris dendritica, 338 Cladoniae, 95 Cladosporium astericola, 337; herbarum, 240 Clarkeinda, 61 Clathrus locellus, 316 Clavaria, 178; abietina, 178; botrytis, 178; corniculata, 178; cristata, 178; mucida, 178; subcaespitosa; 231; vermicularis, 178 Cleavage in the sporangia of certain fungi, Method of, 143 Clitocybe illudens, 43; odora, 187; phyllophila, 26; subhirta, 26 Coleosanthus, 106, 107; grandiflorus,

Coleosporium, 114, 244-254, 297, 299,

302, 308; arnicale, 113-115; carneum, 250, 252, 308; delicatulum, 254-

257; elephantopodis, 252; Helianthi, 244-253; inconspicuum, 244, 245, 247-253; Ipomoeae, 252, 297, 299-301; Laciniariae, 253; Madiae, 114, 115; minutum, 252; ribicola, 299-302; Solidaginis, 250, 252, 256, 302, 306-310; Terebinthinaceae, 303, 250-253 Coleosporium - I, Notes on some species of, 244; - II, 297 Colletotrichum Boehmeriae, 88 Collybia butyracea, 93; cirrata, 93; clusilis, 93; conigena, 93; distorta, 93; dryophila, 93; Hariolorum, 93; lilacina, 93; nummularia, 93; semitalis, 93; strictipes, 93; subdryophila, 93; tuberosa, 93; velutipes, Coniophora cerebella, 179 Contributions to our knowledge of Oregon fungi - I, 173 Convolvulus, 299; arvensis, 298; repens, 298; sepium, 298, 305 Cookeina Tricholoma, 338 Coprinus, 61, 220, 268, 271, 317, 318; atramentarius, 135; comatus, 135, 136; micaceus, 44; Spraguei, 274 Coreopsis, 109, 110, 244, 245, 247, 248, 250, 251; delphinifolia, 251; lanceolata, 248, 251; major, 246-248, 250, 251, 298, 304; major Oemleri, 251; major rigida, 251; mexicana, 109, 110; mutica, 110; tinctoria, 303; tripteris, 246, 248, 251; verticillata, 246-248, 250-252, 298, 303-305 Coriolopsis Copelandii, 93 Coriolus sericeohirsutus, 44 Cornus Nuttallii, 175 Corticium lactescens, 178; salmonicolor, 91 Cortinarius, 44; semisanguineus, 45 Corylus californica, 174, 183 Costantinella, 103; cristata, 103 Craterellus cornucopioides, 178 Crucibulum vulgare, 198 Cryptoporus volvatus, 181 Cryptosporella anomala, 46 Cultures of heteroecious rusts, 1920-1921, 228 Cyathus stercoreus, 198; vernicosus, Cydonia japonica, 285, 294; sinensis, 285; vulgaris, 282, 284, 285, 294 Cylindrophora, 337 Cylindrosporium acerinum, 198 Cyphella Heveae, 91; marginata, 179 Cystopus, 146 Dacryomyces aurantia, 177

Eriophorum, 119 Daldinia vernicosa, 174 Dark-spored agarics - I, 61; - II, Eupatorium, 107, 108; atriplicifolium, 121; —III, 200; — IV, 258 107; maculatum, 254; pansamalense, Dasyspora conferta, 118; Millefolii, 108; urticaefolium, 108 Eurotiopsis, 242; Gayoni, 241 118; Synedrellae, 117 Davis, W. H., Urocystis Agropyri on Eurotium, 241 Euthamia, 254-256; caroliniana, 255redtop, 279 Deconica, 258, 268; atrorufa, 263; 257; graminifolia, 254-257, 306; bryophila, 261; bulbosa, 260; bullaleptophylla, 257 cea, 258, 262, 263; coprophila, 259; Excidia recisa, 177; Zelleri, 177 nuciseda, 259; polytrichophila, 261; pyrispora, 261; rhomboidospora, Fagus, 1, 2, 5 259; scatigena, 258; semistriata, Farlowianae, Reliquiae, 99 260; subviscida, 262; tomentosa, Fink, B., An addition to the distribu-259 tion of a rare fungus, 49; A new Delitescor, 258 lichen from an unusual substratum, Dendrostilbella, 241, 242; Boydii, 242 95 Fitzpatrick, H. M., Thomas, H. E., and Kirby, R. S., The Ophiobolus causing take-all of wheat, 30 Derbesia, 148 Diagnoses of American porias - I, 1 Diaporthe Menispermi, 233; Triostei, 233 Flaveria angustifolia, 118; campes-Diatrype, 174 tris, 118 Diatrypella, 100; ciliatula, 100 Fomes annosus, 181; applanatus, 181; Dicaeoma Balsamorrhizae, 112; Brickconnatus, 9; igniarius, 94, 181; elliae, 106; cubense, 116, 117; lamaensis, 93; leucophaeus, 181; Eriophorii, 119; Hemizoniae, 115; lignosus, 91; lineatus, 92; pinicola, Kuhniae, 106; nudum, 115; Sar-181; populinus, 9; pseudoferreus, cobati, 229; Synedrellae, 116; trian-91; roseo-albus, 92, 93; roseus, 182, nulatum, 111 183; semilaccatus, 92 Didymella Agrostidis, 233 Fragrant polypore, A, 46 Diervilla Lonicera, 101 Fraser, W. P., Cultures of heteroe-Dimorphotheca, 119, 120; cuneata, cious rusts, 1920-1921, 228 IIQ Fungi - I, Contributions to Discina ancilis, 175 knowledge of Oregon, 173 Distichlis stricta, 229, 230 Fungi, Illustrations of, 25 Distribution of a rare fungus, An Fungi, Index to illustrations of, 332 addition to the, 49 Fungi, New Japanese, 81, 282 Dodecatheon pauciflorum, 229, 230 Fungus, An addition to the distribu-Drosophila, 61, 121, 138, 139, 150, tion of a rare, 49 264; appendiculata, 63, 64, 71, 141, Fusarium, 337; oxysporum, 51 265; atrofolia, 71; delineata, 68; echiniceps, 68, 69; fragilis, 64; Galerula Hypni, 27 hololanigera, 70; hydrophila, 72; Ganoderma, 92; Curranii, 92; leptolacrymabunda, 67, 69; madeodisca, pum, 92; oregonense, 43, 92; um-62, 65, 72; Peckiana, 68; pecosense, braculum, 92 63; rigidipes, 70; Storea, 66, 72 Gaultheria Shallon, 178 Gautieria, 196; morchelliformis, 196; Echinodontium tinctorum, 180 Parksiana, 196, 197 Electra, 110; Galeottii, 110 Geaster, 311, 316, 321; delicatus, 191; Elephantopus, 250; carolinianus, 248hygrometricus, 192; hygrometricus 250, 298, 303, 304; nudatus, 298; giganteus, 192; lagenaeformis, 192; tomentosus, 249, 254, 298, 303-306 Eleutheranthera ruderalis, 116, 117 Morganii, 192; saccatus, 192 Gelatinosporium abietinum, 101 Elliott. J. A., Some characters of the Genus of gill-fungi, Another greensouthern tuckahoe, 222 Elymus, 279; arenarius, 279; canadenspored, 96 Geophila, 125; capnoides, 74 sis, 279; robustus, 279; virginicus,

Emilia sagittata, 119; sonchifolia, 119

Equisetum, 175

Geopyxis cupularis, 175

nus of, 96

Gill-fungi, Another green-spored ge-

our

Notes on some species of Coleo-Glaux, 229; maritima, 228-230; Michsporium - I, 244; - II, 297 auxiana, 228 Helianthus, 244, 245, 247, 250; angus-Gleditsia, 2 tifolius, 298, 304; annuus, 255, 304; Gliocladium penicillioides, 101 Gloeosporium acutilobum, 233; Balsameae, 337; bicolor, 337; limetticola, 335 Glomerella, 84; Cinnamomi, 82 Gnomonia Coryli, 174 Gnomoniopsis acerophila, 233 Godronia Nemopanthis, 101; turbinata, 101 Gomphidius, 121, 200; flavipes, 124; furcatus, 125; glutinosus, 44, 122; jamaicensis, 121; maculatus, 124, 125; nigricans, 123; oregonensis, 121, 191; tomentosus, 121; vini-304 color, 121, 123; viscidus, 44, 122, Oryzae, 81, 82 Green-spored genus of gill-fungi, Another, 96 Hemizonella, 114, 115 Grossularia, 301; Cynosbati, 301; di-Hemizonia, 114, 115 varicata, 300, 302; hirtella, 300, 302; inermis, 300-302; innominata, 300, 302; leptantha, 300, 301; missouriensis, 300, 302; rhodospiloides, 236, 237 300-302; setosa, 301 Guignardia Cinnamomi, 84 Heteromeles, 196 Gymnochilus roseolus, 268 Hevea brasiliensis, 91 Gymnomyces Gardneri, 198; pallidus, 197, 198 Gymnopilus flavidellus, 28; polysketch of the, 311 chrous, 75, 141, 142 Gymnopus velutipes, 45 Gymnosporangium, 283-286, 290, 291, 293; alpinum, 294; amelanchieris, 289; asiaticum, 282-285, 287, 292, 294; blasdaleanum, 295; chinensis, 287, 293, 294; clavariaeforme, 295; clavariiforme, 287, 289, 293; claviaeforme, 287, 293; clavipes, 289; confusum, 284, 293, 294; Haraea-Hydrodictyon, 149 num, 283, 284, 287, 290, 294; hemisphaericum, 289, 294; Idetae, 287, 294; japonicum, 283-285, 288, 293, 294; Juniperi, 294; juniperinum, 289, 294; koreaense, 294; koreaensis, 284, 287; Miyabei, 288, 294; Photiniae, 284, 285, 293; Sabinae, 295; Shiraianum, 291, 294; spiniferum, 284, 294; tremelloides, 288, 294; Yamadae, 285, 287, 293; Yamadai, 286, 293 Gyrocephalus rufus, 177 Gyromitra esculenta, 176 Harpaecarpus, 115

Hebeloma, 75; luteum, 29

Hedgoock, G. G., and Hunt, N. R.,

australis, 250; decapetalus, 246-248, 250, 251, 298, 305; divaricatus, 246-248, 250, 251, 254, 298, 304, 305; doroconoides, 250; Dowellianus, 246, 248; Eggertii, 250; giganteus, 246, 247, 250; glaucus, 246-248, 250, 251; grosseserratus, 250; hirsutus, 246-248, 250; microcephalus, 246-248, 250, 251; occidentalis, 248, 250, 254, 303, 304; Radula, 246, 247, 251, 298; saxicola, 251; strumosus, 251; tuberosus, 248, 251, Helminthosporium macrocarpum, 82; Helvella crispa, 176, 198; elastica, 176; infula, 176; lacunosa, 176 Herpotrichia, 236, 237; albidostoma. 235, 236, 238; diffusa, 236-238; incisa, 238; rhodomphala, 237; Heteroecious rusts, Cultures of, 228 Hexagonia carbonaria, 183 History of mycological illustration, A Hunt, N. R., and Hedgcock, G. G., Notes on some species of Coleosporium — I, 244; — II, 297 Hydnangium carneum, 335 Hydnotrya ellipsospora, 176 Hydnum auriscalpium, 180; erinaceus, 180; ochraceum, 180; repandum, 180; subfuscum, 180 Hydrocybe ceracea, 44; Earlei, 44 Hygrophorus, 48; caprinus, 48; eburneus, 188; fuligineus, 48; miniatus, 188; miniatus sphagnophilus, 188 Hylocomium triquetrum, 188 Hymenochaete spreta, 179 Hymenopappus carolinensis, 118 Hypholoma, 61, 72, 75, 96, 121, 190; aggregatum, 66; aggregatum sericeum, 66; appendiculatum, 63, 64, 71; Artemisiae, 75; Boughtoni, 67, 68; Candolleanum, 63, 71; capnoides, 72, 74, 191; comaropsis, 75; comatum, 71; confertissimum, 71; coronatum, 71; cutifractum, 64; delineatum, 68; echiniceps, 69; fasciculare, 72-74, 190, 191; flocculentum, 64; fragile, 64; hololanigerum, 70; hydrophilum, 191; lacrymabundum, 66, 67, 191; lateritium, 72; madeodiscum, 62, 63, 71; nitidipes, 75; ornellum, 75; Peckianum, 68; pecosense, 63, 133; perplexum, 73; populinum, 72; Pseudostorea, 66; rigidipes, 70; rugocephalum, 67, 68; subaquilum, 62, 63, 71; sublateritium, 73; velutinum, 191; velutinum leiocephalum, 71; vinosum, 97

Hypoxylon atropunctatum, 174; pruinatum, 51

Illustration, A sketch of the history of mycological, 311
Illustrations of fungi,—XXXIII, 25
Illustrations of fungi, Index to, 332
Index to American mycological literature, 53

Index to illustrations of fungi, XXIII-XXXIII, 332

Inocybe, 9r, 214, 232, 260; fastigiata, 189

Inonotus dryophilus, 44; perplexus, 43

Ipomoea, 299; Batatas, 298; caroliniana, 298, 299; fistulosa, 303; lacunosa, 298, 299; pandurata, 298, 299, 303; triloba, 298, 299 Iris foetidissima, 33-35 Irpex, 1; lacteus, 180

Jackson, H. S., New or noteworthy rusts on Carduaceae, 104
Japanese fungi, New, 81, 282
Juniperus chinensis, 282, 283, 285, 286, 290, 293, 294; chinensis procumbens, 282, 286, 293, 294; littoralis, 292-294; nana, 294, 295; nipponica, 294; rigida, 285, 288, 293, 294

Justicia procumbens, 87

Kuhnia, 107

Kirby, R. S., Fitzpatrick, H. M., and Thomas, H. E., The Ophiobolus causing take-all of wheat, 30 Knowledge of Oregon fungi — I, Contributions to our, 173 Krieger, L. C. C., A sketch of the history of mycological illustration (higher fungi), 311; Schizophyllum commune with a stipe, 47

Laccaria laccata, 44 Laciniaria acidota, 303; elegans, 246, 248, 298; elegantula, 298; gracilis, 298; graminifolia, 252, 298; pycnostachya, 298; tenuifolia, 298 Lactarius pyrogalus, 188 Laetiporus sulphureus, 44 Lagenidium entophytum, 145 Lagophylla, 115 Lapsona communis, 176 Lasiosphaeria strigosa, 174 Lentinula detonsa, 338 Lentinus, 223 Lentodiellum concavum, 338 Lenzites saepiaria, 93, 188; vialis, 189 Lepiota, 96, 186, 214, 215; amianthina, 214; clypeolaria, 186; cretacea, 44; granulosa, 186; Morgani, 96; naucina, 221; procera, 43, 315; pulcherrima, 186; rubrotincta, 186; rubrotinctoides, 186 Leptosphaeria Collinsoniae, 233 Leptothyrium conspicuum, 233 Lichen from an unusual substratum, A new, 95 Licipenicillium insigne, 101 Life history of an undescribed ascomycete isolated from a granular mycetoma of man, 239 Lilliputia Gaillardii, 101 List of the myxomycetes of the Cayuga Lake Basin, 38 Lophodermium Thuyae, 337 Lycoperdon, 321; cruciatum, 192; elegans, 192; fuscum, 192; gemmatum, 192; pratense, 192; pusillum, 192; pyriforme, 192; rimulatum, 192; solidum, 320; Wrightii, 193

Macbride's North American slimemoulds, 233 Macrosporium, 44, 337 Madaria, 115 Madia, 114, 115; Nuttallii, 115 Marasmius plicatulus, 188 Meeting, The Toronto, 50 Melampodium divaricatum, 115, 117 Melanconium oblongum, 46 Melanoleuca eduriformis, 27; subannulata, 187; Thompsoniana, 26 Melanotus, 61 Meliola Abietis, 174 Merulius brassicaefolius, 184; pilosus, 184, 185; rugulosus, 185; tremulosus, 185 Method of cleavage in the sporangia of certain fungi, 143 Microdiplodia Populi, 232 Micropuccinia, 108, 109, 118, 119; conferta, 119; Millefolii, 119; Synedrellae, 117 Miles, L. E., A new species of Myriangium on pecan, 77 Monascus purpureus, 42

Monochytrium, 148 175; Morchella, 316; angusticeps, crassipes, 175, 316; deliciosa, 175; esculenta, 315, 316; semilibera, 176 Mucor, 143, 144; mucedo, 144, 146, 163, 164, 166; racemosus, 161; spinescens, 149, 161 Muenscher, W. C., and Wann, F. B., A preliminary list of the myxomycetes of the Cayuga Lake Basin, 38 Murrill, W. A., A fragrant polypore, 46: Another green-spored genus of gill-fungi, 96; Dark-spored agarics -I, 61; —II, 121; —III, 200; — IV, 258; Hygrophorus caprinus, 48; Illustrations of fungi - XXXIII, 25; Index to illustrations of fungi, XXIII-XXXIII, 332; The Toronto meeting, 50 Mycena palustris, 93; pura, 44 Mycetoma of man, Ascomycete isolated from a granular, 239 Mycogone puccinioides, 198; roseola, 198 Mycological illustration, A sketch of the history of, 311 Arundinariae, Mycosphaerella, 232; 86; bambusifolia, 85 Myriangium, 80; tuberculans, 80 Myriangium on pecan, New species of, 77 Myxomycetes of the Cayuga Lake Basin, 38 Naematelia encephala, 177

303, 304 Naucoria, 75; pennsylvanica, on, 77 semiorbicularis, 189 Nectria, 174; coccinea, 174, 175; Coryli, 174; episphaeria, 174; galligena, 174 Neopeckia, 236, 237; diffusa, 235-237 Neurolaena lobata, 119 Neurophyllum viride, 96 New Japanese fungi, Notes and translations - XI, 81; - XII, 282 nea, 101 New lichen from an unusual substratum, 95 New or noteworthy rusts on Carduaceae, 104 New Septobasidium on Pinus Strobus, New species of Myriangium on pe-303, 304 can, 77 North American slime-moulds, Macbride's, 233 Notes and brief articles, 42, 90, 231,

Notes on some species of Coleospo-

rium — I, 244; — II, 297

Noteworthy rusts on Carduaceae, 104 Nyssa, 336 Odontia fragilis, 180 Olpidiopsis, 146, 147, 150-152; major, 151; Saprolegniae, 152, 154, 155 Olpidium Dicksonii, 148, 151; Viciae, 152 Oospora, 337 Ophiobolus, 30-32, 35; cariceti, 35, 36; eucryptus, 32; graminis, 30-33, 35, 36; herpotrichus, 30, 36 Ophiobolus causing take-all of wheat, Oregon fungi - I, Contributions to our knowledge of, 173 Oryza sativa, 81 Ostrya, 5 Otidea leporina, 175 Overholts, L. O., Diagnoses of American porias - I, 1 Oxalis oregana, 198 Pachyma Cocos, 92, 222, 223 Panaeolus, 130, 136; campanulatus, 275; epimyces, 135 Panicum, 198; proliferum, 89; proliferum acuminatum, 89 Parthenice mollis, 108 Parthenium, 252; argentatum, 109; Hysterophorus, 109, 116; incanum, 109; integrifolium, 246, 252, 253, Paxillus pubescens, 122 Pecan, A new species of Myriangium Penicillium insigne, 101 Peniophora cinerea, 179; glabra, 179; glebulosa, 179; incarnata, 179 Peridermium, 302; acicolum, 302, 310; delicatulum, 254; inconspicuum, 244; montanum, 302, 310 Pezicula pruinosa, 102 Peziza, 102; Acetabulum, 317; quer-Phacidium Balsameae, 337; expansum, 337; planum, 337 Phaeosphaeria Bambusae, 86; Oryzae, Pharbitis, 299; barbigera, 298, 299; hederacea, 298, 299; purpurea, 298, Phialea alniella, 175 Pholiota, 42, 141, 214, 220; duroides, 75; Howeana, 141; Johnsoniana, 141; ornella, 75; spectabilis, 189 Phomopsis Impatientis, 233

Photinia, 284; laevis, 293; villosa, 283,

284, 293, 295

Phragmidium, 317 Pleurotus, 187, 232; ostreatus, 44; sapidus, 188; serotinus, 188 Phycomyces, 149, 166; nitens, 149, Pluteus, 76, 98 158, 166 Phyllostachys bambusoides, 85; pu-Polyphagus Euglenae, 147 Polypore, A fragrant, 46 berula, 85 Phyllosticta, 86, 233 Polyporus anebus, 92; arcularius, 182; Physoderma Maydis, 84; Zeae-Maybasilapiloides, 43; Bracei, Broomei, 10; caliginosus, 92; cardis, 85 bonarius, 91; chioneus, 182; cuti-Phytophthora, 146, 336; Faberi, 91; fractus, 91; dryadeus, 182; fastuinfestans, 148, 336 Piggotia Vaccinii, 337 osus, 92; floriformis, 182; fragilis, Pilobolus, 148, 149, 158, 159, 162, 182; galactinus, 182; hirsutulus, 166, 167; crystallinus, 148, 160, 182; hirtus, 182; lucidus, 93; Macounii, 182; Mariannus, 92; minor-161, 163-168 Pilosace, 61, 72, 75, 76, 96, 121, 136; Mylittae, 43; Mylittae, 43; pargamenus, 3, 4; picipes castaneus, 182; algeriensis, 76, 136; eximia, 76; pubescens, 182; rhodophoeus, 92; Peckii, 97 rigidus, 9, 11; rufescens, 183; sen-Pinus, 250-252, 256, 301; Banksiana, sibilis, 183; Spraguei, 183; squamo-250; Bungeana, 300; canariensis, sus 315; trabeus, 182; tulipiferus, 256; caribaea, 247, 255-257, 299, 4; tumulosus, 43; versicolor, 182; 300, 306, 307, 309; clausa, 255; volvatus, 94; zonatus, 183 contorta, 255-257, 302, 305, 306-Polysaccum, 321; turgidum, 194 309; Coulteri, 255-257, 306, 307. Polythelis Thalictri, 177 309; densiflora, 255; edulis, 247, Polytrichum, 261 255, 256, 300-302, 306; echinata, Populus, 5, 181, 184; grandidentata, 248, 250-252, 255-257, 297, 299, 43, 47 303, 306-309; Girardiana, 300; Poria, 1, 183; ambigua, 1, 2, 4; Broomglabra, 247, 248, 255-257; heteroei, 11; contigua, 183; ferruginosa, phylla, 309; Lambertiana, 56; May-5-7; Macouni, 6; marginella, 7; niriana, 255-257, 300, 306; monogrescens, 7, 11; odora, 10, 11; sanphylla, 255, 300; montana, 255, 306; guinolenta, 183; tenuis, 6; undata, monticola, 56, 58, 59; nigra, 307; 10; viticola, 6, 7 nigra austriaca, 309; nigra poireti-Porias, Diagnoses of American, - I, 1 ana, 256; palustris, 248, 251, 252, Poronidulus conchifer, 51 254-257, 298, 306, 307; Pinea, 300, Pourthiaea villosa, 284 302; pungens, 304, 306-308; radiata, Pratella, 200 247, 305, 307, 309; resinosa, 255, 256, Preliminary list of the myxomycetes 305, 307; rigida, 252, 254-257, 300, of the Cayuga Lake Basin, 38 Protococcus, 95 303, 306-309; scopulorum, 255-257, 306, 308, 309; serotina, 252, Protomyces bellides, 147 255-257, 300, 306; strobiformis, Prunulus, 266 300; Strobus, 58, 300; taeda, 252, Prunus, 5; demissa, 91 255-257, 298, 300, 304, 306-309; Psalliota, 200 Thunbergii, 255; virginiana, 245-Psathura, 264 248, 250, 251, 300 Psathyra, 96, 264, 269, 278; australis, 278; gyroflexa, 274; microsperma, Pirus, 295; Aria kamaoensis, 294; 264; multipedata, 265; obtusata, Aucuparia, 293, 294; Aucuparia japonica, 294; communis, 1, 88, 268; persimplex, 268; pholidota. 285, 294; Malus, 286, 290, 293; polytrichophila, 261, 268; Miyabei, 294; spectabilis, 286, 287, prunuliformis, 278; roseola, 268; 293; sinensis, 88, 282, 284, 290, semivestita, 265, 269; silvatica, 269; 291, 293, 294; Toringo, 285-287, striatula, 278; umbonata, 267; ves-290, 293; Zumi, 289, 290, 294 tita, 265, 269 Plantago, 230; eriopoda, 230 Psathyrella, 258, 264, 269, 271; an-Plasmopara Wildemaniana macrospogusticeps, 275; atomata, 272, 275; Bartholomaei, 272, 273; betulina, ra, 87 Pleomassaria Magnoliae, 100; maxi-273; castaneicolor, 272; Clementsii, ma, 100 274; crenata, 278; debilis, 274, 278; disseminata, 270; distantifolia, Pleonectria berolinensis, 175

277; falcifolia, 278; gracilis, 278; gracillima, 274; graciloides, 276, 277; hiascens, 278; hirta, 272; leucostigma, 271; minima, 272; minutula, 269, 270; odorata, 276; petasiformis, 276; rupincola, 278; sulcata, 274; tenera, 271 Pseudolpidium Aphanomycis, 148. 151, 154, 156, 158 Pseudoplectania melaena, 175 Pseudospora Lindstedii, 148 Pseudotsuga, 176, 177, 181, 196; taxifolia, 43, 176, 178-181, 184, 186, 196 Psilocybe, 138, 258, 263, 264; atomatoides, 277; conissans, 72 Puccinia, 104-106, 108, 114, 115, 120; abrupta, 113; Actinellae, 118; additicia, 109, 110; artemisiicola. 118; Asteris, 118; aut-eu-Puccinia, 105; Balsamorrhizae, 111, 112; Barroetiae, 106; brachy-Puccinia, 106, 111, 112; Brickelliae, 106; Burnettii, 176; clara, 104; cognata, 113; conferta, 118; Coreopsidis, 109, 110; Diaziana, 116, 117; Distichlidis, 228-230; Electrae, 110; Eleutherantherae, 116, 117; Emiliae, 116, 119, 120; Eriophorii, 119; Eupatorii, 107; ferox, 113; Flaveriae, 117; gigantispora, 176; Grindeliae, 118; Hemizoniae, 114. 115; hyalina, 104; invelata, 113; irregularis, 113; Kuhniae, 106, 107; Lagophyllae, 115; Lapsanae, 176; Madiae, 114, 115; Melampodii, 115, 116; Millefolii, 118; mirifica, 111; nuda, 113-115; pachyspora, 107; Parthenices, 108; Parthenii, 109; praealta, 105; recedens, 119; recondita, 118; redempta, 107; Silphii, 116; solida, 116, 117; spilanthicola, 116; subdecora, 105; subnitens, 228-230; Synedrellae, 115, 116; Tetranthi, 116, 117; tolimensis, 108; Tonduziana, 104, 105; triannulata, III; Tridacis, II6, II7; turgidipes, II0; vaga, II2, 113; Verbesinae, 113; Wyethiae, 111, 112; Zinniae, 116, 117 Pucciniastrum, 46; americanum, 46; arcticum, 46 Pyropolyporus Williamsii, 92 Pythium entophytum, 145; intermedium, 156, 158; proliferum, 157

Quamoclit coccinea, 298, 299; Quamoclit, 298 Quercus, 2, 5, 176; Garryana, 174, 175, 183, 195, 196, 198, 199

Radulum Owensii, 180 Ramularia cilinodis, 337; Eamesii, 233; minax, 337; Oxalidis, 198 Ranunculus apetalus, 229, 230 Redtop, Urocystis Agropyri on, 279 Reliquiae Farlowianae, 99 Rhaphidophora graminis, 36 Rhizopogon diplophloeus, 195; induratus, 195; maculatus, 196, 335; roseolus, 196; rubescens, 196 Rhizopus, 143, 144, 337; nigricans, 143, 149, 158-160, 163-166, 168 Rhodochytrium, 147 Ribes, 175, 301; alpinum, 300; americanum, 300-302; aureum, 300-302; coloradense, 301; fasciculatum, 300, 302; inebrians, 300-302; malvaceum, 300; mescalerium, 301; montigenum, 301; nigrum, 254, 300, 302; odoratum, 300-302; pumilum, 301; sanguineum, 301; vulgare, 300, 302; Wolfii, 301 Robinia Pseudacacia, 1 Roestelia, 282, 283; cancellata, 295; koreaensis, 282-284, 294; Photiniae, 284, 293; Pourthiaeae, 293; solenoides, 294; solitaria, 294 Roskovites granulatus, 45 Rozella, 146, 147, 150, 151 Rubus, 190; idaeus aculeatissimus, 46; occidentalis, 46; pubescens, 46; strigosus, 46; triflorus, 46 Rudbeckia laciniata, 246 Russula, 45; emetica, 45; foetens, 315; nigricans, 315; ochrophylla, 188; virescens, 315 Rusts, Cultures of heteroecious, 228 Rusts on Carduaceae, New or noteworthy, 104

Saccardinula Alni, 233 Saccharum Narenga, 87 Salix, 5, 174, 175; alba, 47 Sambucus glaucus, 177; pubens, 277 Sanicula Menziesii, 174 Saprolegnia, 146-148, 150-152, 154, 155, 158-160, 163, 166 Sarcoscypha coccinea, 175 Sasa paniculata, 86 Schizonella melanogrammia, 177 Schizophyllum, 47, 232; alneum. 48; commune, 47, 232; commune stipitatum. 48 Schizophyllum commune with a stipe, Schulzeria, 96 Schwarze, C. A., The method of cleavage in the sporangia of certain fungi, 143 Scirpus maritimus, 229

ticola, 305, 309; multiradiata, 254, Scleroderma, 194, 321; aurantiacum, 298, 303-306, 309; neglecta, 304, 194; cepa, 193, 194; hypogaeum, 193, 194 306, 309; nemoralis, 304; odora, Scleroderris, 102; Cephalanthi, 102 309; petiolaris, 309; pinensis, 309; Pinetorum, 309; pulverentula, 309; Sclerotinia, 91; demissa, 91; gregaria. or pulverula, 309; Purshii, 309; Rid-Seaver, F. J., Macbride's North dellii, 303-306, 309; rigida, 304, slime-moulds, 309; rigidiuscula, 309; rugosa, 254, American 233: Studies in tropical ascomycetes — 255, 298, 303, 304, 306, 309; rupes-I, 235 tris, 309; sempervirens, 304, 306; Seaver, Fred J., Uredinales collected serotina, 303, 304, 306, 309; speciby, 12 osa, 254, 255, 303, 304, 306, 309; Secotium tenuipes, 335 squarrosa, 303-306, 309; stricta, 309; tortifolia, 304, 309; unigulata, Senecio, 119, 120; aureus, 254, 303, 305; Cineraria, 120; obovatus, 309; Vaseyii, 309 Some characters of the southern 303, 305 tuckahoe, 222 Sepedonium chrysospermum, 198 Sorbus, 295; Aucuparia, 293; japoni-Septobasidium, 55-57; pinicola, 55-58; ca, 293, 294; sambucifolia pseudopseudopedicillatum, 56 gracilis, 294 Septobasidium on Pinus Strobus, A new, 55 Sparassis radicata, 177 Spartina Michauxiana, 228; pectinata, Septocladia dichotoma, 231 Septoria Acetosella, 233; Coreopsidis, 228, 230 337; hieracicola, 233 Spathularia clavata, 176 Shear, C. L., Life history of an Species of Coleosporium - I, Notes on some, 244; — II, 297 Species of Myriangium on pecan, ascomycete isolated undescribed from a granular mycetoma of man, A new, 77 Sphaeria, 102; albidostoma, 238; ca-Sidalcea, 176 Silphium, 116, 250, 252; angustatum, riceti, 32-36; ciliatula, 100; diffusa, 253; asperrimum, 253; asteriscum, 237; eucrypta, 32-35; rhodomphala, 246, 250, 252, 253; compositum, 237 253; dentatum, 253; glabrum, 253; Sphaerobolus stellatus, 198 Sphaeronema pruinosum, 102 gracile, 253; integrifolium, 246, 248, 250, 252, 253; laciniatum, 253; laevigatum, 253; perfoliatum, Sphaeropsis ulmicola, 335 Sphaerulina pallens, 337 246; pinnatifidum, 253; scaberri-Sphagnum cuspidatum, 103; Torreymum, 253; Simpsonii, 298; terebinanum, 103 Spilanthes oleracea, 116 thinaceum, 253, 298; trifoliatum, 246, 252, 253 Sporangia of certain fungi, Method Sketch of the history of mycological of cleavage in the, 143 Sporidesmium Fusus, 100 illustration (higher fungi), 311 Sporidinia, 146, 148, 149, 159; gran-Slime-moulds. Macbride's North dis, 148, 158-164, 166-168 American, 233 Snell, W. H., A new Septobasidium Stagonospora Meliloti, 233; tetrameon Pinus Strobus, 55 ra, 337 Stamnaria Persoonii, 175 Solidago, 250, 302-305, 307-309; am-Stemphylium, 44 plexicaulis, 309; austrina, 309; bi-Stereum fuscum, 178; purpureum, color, 249, 254, 303-307, 309; Bootii, 309; brachyphylla, 309; cae-178 Stichospora Madiae, 115 sia, 303, 304; canadensis, 247, 254, Stieronema ciliatum, 228 298, 303-306, 309; celtidifolia. 309; Chandonnetii, 309; Chapmannii, 303, Stilbum Rhois, 103 304, 306, 309; concinna, 309; Cur-Stropharia, 76, 121, 125, 136, 200; acuminata, 128; adnata, 133; aetissii, 309; decumbens, 309; Drumruginosa, 128, 140, 142, 190; albomondii. 309; erecta, 254, 303, 306, 309; fistulosa, 298, 303, 304, 306, cyanea, 140; albonitens, 140; am-

309; glomerata, 309; hispida, 303, 306, 309; juncea, 249, 254, 298,

303-306, 309; lancifolia, 309; mon-

bigua, 190; annellariformis, 129;

bilamellata, 130, 131, 137, 211; caesifolia, 136, 137; caesiospora,

134; campestris, 136; coprinophila, 135; coronilla, 131, 137, 139, 141; cotonea, 66; depilata, 134; distans, 126; **elegāns**, 140; epimyces, 76, 135; Feildeni, 141; Hardii, 134, 135; Howeana, 141; irregularis, 65, 141; Johnsoniana, 141; mammillata, 129; melanosperma, 130, 131; merdaria, 138; micropoda, 141; obdurata, 131; obturata, 141; paradoxa, 129; pseudocyanea, 140; rugomarrugoso-annulata, ginata, 190; 139; Schraderi, 141; semiglobata, 132-134; siccipes, 132; siccipes radicata, 132; squamosa, 126, 127; stercoraria, 133; subbadia, 139; submerdaria, 138; tenuis, 137; umbilicata, 142; umbonatescens, 128 Strumella coryneoidea, 199 Studies in tropical ascomycetes - I, Substratum, A new lichen from an unusual, 95 Syncephalastrum, 147; cinereum. 149; racemosum, 149 Syncephalis, 147; cordata, 147; nodosa, 147; pycnosperma, 147; Wynneae, 147 Synchytrium, 151, 167; decipiens. 148, 155, 156, 158, 160, 163, 167; endobioticum, 149; Puerariae, 147, 156, 158; pulvereum, 337; Succisae, 149; Taraxaci, 147 Synedrella nodiflora, 115 Take-all of wheat, The Ophiobolus causing, 30 Tanaka, T., New Japanese fungi. Notes and translations - XI, 81; -XII, 282 Tetranthus hirsutus, 116 Thalictrum occidentale, 177 Thaxter, R., Reliquiae Farlowianae, Thelephora palmata, 178 Thelocarpon, 95; fimicola, 96 Thomas, H. E., Fitzpatrick, H. M., and Kirby, R. S., The Ophiobolus causing take-all of wheat, 30 Thuja plicata, 179 Thyella, 299 Toronto Meeting, 50 Trametes, 183; carnea, 181, 183; heteromorpha, 184; hispida, 184; suaveolens, 46, 47; tenuis, 6, 183 Tremella frondosa, 177; lutescens, 177; mesenterica, 177; violacea, 177 Trichia, 158 Trichobasis Balsamorrhizae, 112;

Wyethiae, 112 Trichoglossum hirsutum, 176 Tricholoma, 187; bicolor, 187; personatum, 187; subannulata, 187 Tridax procumbens, 116 Trinidad, Uredinales collected by Fred J. Seaver in, 12 Trogia crispa, 188 Tropical ascomycetes — I, Studies in, Tsuga, 181; heterophylla, 179 Tubaria silvatica, 269 Tuber, 176; candidum, 176 Tuckahoe, Some characters of the southern, 222 Tulostoma, 50 Tylopilus alboater, 25 Tylostoma, 50; campestre, 50; verrucosum, 50, 232 Tympanis, 101 Typhula phacorrhiza, 178

Ulmus, 5 Underwoodia, 316 Uredinales collected by Fred J. Seaver in Trinidad, 12 Uredo abdita, 120; Balsamorrhizae, 112; Ipomoeae, 297; Nyssae, 336; Parthenii, 109; ribicola, 299; Terebinthinaceae, 251; triannulata, 111 Urocystis Agropyri, 279, 281 Urocystis Agropyri on redtop, 279 Uromyces Alopecuri, 229, 230; Scirpi, 229; seditiosus, 230; triannulatus, III Urophlyctis pluriannulatus, 174 Urticastrum, 336 Ustilaginoidea Oryzae, 87; Sacchari-Narengae, 87 Ustilago formosana, 89; Panici-proliferi, 89 Ustulina zonata, 91

Vaucheria, 166, 167
Venturia fimbriata, 233
Verbesina, 113, 244, 245, 249, 250; virginica, 247–250, 298, 303, 304
Vernonia, 250; angustifolia, 304; Blodgettii, 248, 304; flaccidifolia, 248, 250, 303, 304; glauca, 248, 249, 298, 303–306; noveboracensis, 247, 248, 254, 303, 304, 306; oligantha, 248; scariosa, 104; triflosculosa, 105
Verticillium pyramidale, 103
Viguiera, 244, 245; deltoidea Parishii, 110

Wann, F. B., and Muenscher, W. C., A preliminary list of the myxomycetes of the Cayuga Lake Basin, 38
Wheat, The Ophiobolus causing takeall of, 30
Woronina, 146, 147, 151
Wyethia, 112

Ximenesia encelioides, 116 Xylaria, 311, 316; Longiana, 174 Zea Mais, 84
Zeller, S. M., Contributions to our knowledge of Oregon fungi — I,
173
Zinnia tenuiflora, 116
Zonanthemus, 115
Zygorhizidium Willei, 148, 151